

Microscopic theory of the interaction of ultracold dense Bose and Fermi gases with electromagnetic field

K.V.Krutitsky ^{*}, F.Burgbacher and J.Audretsch [†]

Fakultät für Physik, Universität Konstanz, Fach M 674, D-78457 Konstanz, Germany
(2d July, 1999)

We present the rigorous microscopic quantum theory of the interaction of ultracold Bose and Fermi gases with the electromagnetic field of vacuum and laser photons. The main attention has been paid to the consistent consideration of dynamical dipole-dipole interactions. The theory developed is shown to be consistent with the general principles of the canonical quantization of electromagnetic field in a medium. Starting from the first principles of QED we have derived the general system of Maxwell-Bloch equations for atomic creation and annihilation operators and the propagation equation for the laser field which can be used for the self-consistent analysis of various linear and nonlinear phenomena in atom optics at high densities of the atomic system. All known equations which are used for the description of the behaviour of an ultracold atomic ensemble in a radiation field can be obtained from our general system of equations in a low-density limit.

I. INTRODUCTION

In the last decade rapid progress is achieved in laser cooling of neutral atoms. This leads to the vigorous development of the new field of atom optics. One can distinguish linear and nonlinear effects in atom optics. Theoretical analysis of the linear effects in atom optics is carried out without consideration of atomic interactions. On the other hand, it was shown that the consideration of dynamical dipole-dipole interactions in the radiation field leads to the nonlinear effects such as a nonlinear Bragg diffraction and the propagation of atomic solitons. That is why in recent years the problem of the interaction of ultracold atoms in the field of external laser radiation has attracted a lot of attention.

The analysis carried out by O.Morice et al. [1] and J.Ruostekoski and J.Javanainen [2,3] was mainly concentrated on the investigation of the properties of the laser radiation modified by atomic dipole-dipole interactions. It was shown that this can be described by the refractive index which is governed by the Clausius-Mossotti relation known from classical optics if we neglect quantum

correlations. If the quantum correlations caused by Bose or Fermi statistics are taken into account, the formula for the refractive index contains additional terms defined by a position dependent correlation function [1–3].

The modification of the properties of the laser radiation should have a back influence on the behavior of ultracold atomic ensemble (for example, the motion of atomic beam). An attempt to consider this back influence was undertaken by several authors [2,4–8]. In order to incorporate two-body interactions G.Lenz, P.Meystre, and E.M.Wright proceeded by eliminating the transverse vacuum field and replacing it by the phenomenological contact potential [4]. Dynamical dipole-dipole interactions which play more important role than contact interactions were taken into account in another papers [2,5–8]. However, in papers by W.Zhang and D.Walls [5] and G.Lenz et al. [6] the averaged polarization of ultracold atomic ensemble was computed as a function of the incident laser field, whereas it should be a function of the macroscopic or the local field which are different from the external laser field due to dynamical dipole-dipole interactions. Wallis [8] paid attention to this fact and used the correct form of the equations for the electromagnetic field. However, his equation for the matter field is written in the lower order of the density in comparison with the equation for the electromagnetic field [9]. This problem was considered also by Y.Castin and K.Mølmer [7] and J.Ruostekoski and J.Javanainen [2]. But the equations which are used in these papers are very complicated because they are written down in terms of the local field and dipole-dipole interactions are presented explicitly in the form of the sum over dipole fields. This makes the analysis very difficult and the equations, which govern the time evolution of the matter fields, remained unsolved.

In our recent paper [9] we have developed the rigorous quantum theory of the interaction of an ultracold atomic ensemble with the electromagnetic field of vacuum and laser photons. The main attention has thereby been paid to the consistent consideration of dynamical dipole-dipole interactions. We have shown that the retardation effects influence significantly the behaviour of atomic ensemble in the radiation field. Making use of the Lorentz-Lorenz relation, we have obtained the general system of equations for atomic creation and annihilation operators and the propagation equation for the laser field which can be used for the self-consistent analysis of various linear and nonlinear phenomena in atom optics at high densities of the atomic system. We have shown that all equations which are used up to now for

^{*}Permanent address: Ulyanovsk Branch of Moscow Institute of Radio Engineering and Electronics of Russian Academy of Sciences, P.B.9868, 48, Goncharov Str., Ulyanovsk 432011, Russia; e-mail: ufire@mv.ru

[†]e-mail: Juergen.Audretsch@uni-konstanz.de

the description of an ultracold atomic ensemble subject to the influence of the laser radiation can be obtained from our general system of equations by series expansion with respect to the density.

In spite of the fact that most of the results obtained in [9] remain valid in the case of Fermi-Dirac gases, we considered explicitly only the case of Bose-Einstein statistics. In addition, our treatment of the electromagnetic field was not consistent with the QED canonical quantization procedure. The vacuum fluctuations of the photon field were separated from the electromagnetic field operator. Such an operation leads to the violation of the QED canonical commutation relations. The aim of the present paper is to fill these gaps.

II. THE MODEL AND HAMILTONIAN

We discuss the combined system of a gas of identical neutral polarizable atoms interacting with an incident monochromatic field $\mathbf{E}_{in}(\mathbf{r}, t)$ with frequency ω_L . The probe field $\mathbf{E}_{in}$ is a classical external field. No operator is related with it. It is fixed and therefore not subject to a dynamics. For example, it may be a standing or running laser field. The quantized center-of-mass motion of the atoms is taken into account. The atoms are represented as point-like neutral two-level atoms with states $|1\rangle$ and $|2\rangle$ with “bare” transition frequency ω_a . The atoms interact via the resonant dipole-dipole interaction of the induced point dipoles. This interaction is mediated by the exchange of *photons* with annihilation and creation operators $\hat{c}_{\mathbf{k}\lambda}$ and $\hat{c}_{\mathbf{k}\lambda}^\dagger$ referring in lowest order to plane waves in vacuum with wave vector $\mathbf{k}$ and polarization λ .

To take into account quantum statistical effects for the two-level atoms, we turn to quantized matter fields represented by two-component vectors

$$\psi(\mathbf{r}, t) = \hat{\psi}_1(\mathbf{r}, t)|1\rangle + \hat{\psi}_2(\mathbf{r}, t)|2\rangle. \quad (1)$$

The equal time commutation relations are

$$\begin{aligned} [\hat{\psi}_i(\mathbf{r}), \hat{\psi}_j(\mathbf{r}')]_q &= [\hat{\psi}_i^\dagger(\mathbf{r}), \hat{\psi}_j^\dagger(\mathbf{r}')]_q = 0, \\ [\hat{\psi}_i(\mathbf{r}), \hat{\psi}_j^\dagger(\mathbf{r}')]_q &= \delta_{ij} \delta(\mathbf{r} - \mathbf{r}'), \\ i, j &= 1, 2, \quad q = c, a, \end{aligned} \quad (2)$$

where $[\hat{A}, \hat{B}]_c$ is the commutator of the operators $\hat{A}$ and $\hat{B}$, and $[\hat{A}, \hat{B}]_a$ is the anticommutator, which corresponds to Bose-Einstein or Fermi-Dirac statistics, respectively. In terms of the operators $\hat{\psi}_1$ and $\hat{\psi}_2$ and in the multipolar formulation of QED [10,9] the Hamiltonian operator takes then the form:

$$\begin{aligned} \hat{\mathcal{H}} &= \hat{\mathcal{H}}_A + \hat{H}_F + \hat{\mathcal{H}}_{AI} + \hat{\mathcal{H}}_{AF}, \\ \hat{\mathcal{H}}_A &= \sum_{j=1}^2 \int d\mathbf{r} \hat{\psi}_j^\dagger(\mathbf{r}, t) \left(-\frac{\hbar^2 \nabla^2}{2m} \right) \hat{\psi}_j(\mathbf{r}, t) \end{aligned} \quad (3)$$

$$\begin{aligned} &+ \int d\mathbf{r} \hat{\psi}_2^\dagger(\mathbf{r}, t) \hbar \omega_a \hat{\psi}_2(\mathbf{r}, t), \\ \hat{H}_F &= \sum_{\mathbf{k}\lambda} \hbar \omega_k \hat{c}_{\mathbf{k}\lambda}^\dagger(t) \hat{c}_{\mathbf{k}\lambda}(t), \\ \hat{\mathcal{H}}_{AI} &= - \int d\mathbf{r} \hat{\mathbf{P}}(\mathbf{r}, t) \mathbf{E}_{in}(\mathbf{r}, t), \\ \hat{\mathcal{H}}_{AF} &= - \int d\mathbf{r} \hat{\mathbf{P}}(\mathbf{r}, t) \hat{\mathbf{D}}_{mic}(\mathbf{r}, t). \end{aligned}$$

Here we have introduced the polarization operator $\hat{\mathbf{P}}(\mathbf{r}, t)$ of the ultracold ensemble

$$\hat{\mathbf{P}}(\mathbf{r}, t) = d \hat{\psi}_1^\dagger(\mathbf{r}, t) \hat{\psi}_2(\mathbf{r}, t) + H.c. = \hat{\mathbf{P}}^+(\mathbf{r}, t) + \hat{\mathbf{P}}^-(\mathbf{r}, t), \quad (4)$$

and the operator of the microscopic displacement field is given by

$$\hat{\mathbf{D}}_{mic}(\mathbf{r}, t) = \sum_{\mathbf{k}\lambda} i \sqrt{\frac{2\pi\hbar\omega_k}{V}} \mathbf{e}_\lambda \hat{c}_{\mathbf{k}\lambda} \exp(i\mathbf{k}\mathbf{r}) + H.c. \quad (5)$$

We assume that all contact interactions can be neglected. In order to find the conditions under which this approximation is valid, it is necessary to estimate the ratio U_d/U_g , where U_d and U_g are the mean energies of the dipole-dipole interaction and the ground-state collisions, respectively. Treating the ground-state collisions in terms of s-wave scattering and restricting to dipole optical transitions we get the following inequality [8,9]

$$\frac{U_d}{U_g} \gg 37.5s, \quad (6)$$

where s is the saturation parameter of the atomic transition. Thus, the ground-state collisions are negligible if the saturation parameter s is of the order of 0.01 or higher. This shows, that we don't need very high values of the parameter s in order to neglect contact interactions.

III. HEISENBERG EQUATIONS OF MOTION FOR THE ATOMIC AND PHOTONIC OPERATORS

Making use of the Hamiltonian (3) and the commutation relations (2) we obtain the following Heisenberg equations of motion for the atomic and photonic operators:

$$\begin{aligned} i\hbar \frac{\partial \hat{\psi}_1(\mathbf{r}, t)}{\partial t} &= -\frac{\hbar^2 \nabla^2}{2m} \hat{\psi}_1(\mathbf{r}, t) - d \mathbf{E}_{in}^-(\mathbf{r}, t) \hat{\psi}_2(\mathbf{r}, t) \\ &- \hat{\psi}_2(\mathbf{r}, t) d \mathbf{E}_{in}^+(\mathbf{r}, t) \\ &- \hbar \sum_{\mathbf{k}\lambda} g_{\mathbf{k}\lambda}^* \hat{c}_{\mathbf{k}\lambda}^\dagger(t) \exp(-i\mathbf{k}\mathbf{r}) \hat{\psi}_2(\mathbf{r}, t) \end{aligned}$$

$$- \hbar \hat{\psi}_2(\mathbf{r}, t) \sum_{\mathbf{k}\lambda} g_{\mathbf{k}\lambda} \exp(i\mathbf{k}\mathbf{r}) \hat{c}_{\mathbf{k}\lambda}(t), \quad (7)$$

$$\begin{aligned} i\hbar \frac{\partial \hat{\psi}_2(\mathbf{r}, t)}{\partial t} = & -\frac{\hbar^2 \nabla^2}{2m} \hat{\psi}_2(\mathbf{r}, t) + \hbar \omega_a \hat{\psi}_2(\mathbf{r}, t) \\ & - \mathbf{d} \mathbf{E}_{in}^-(\mathbf{r}, t) \hat{\psi}_1(\mathbf{r}, t) - \hat{\psi}_1(\mathbf{r}, t) \mathbf{d} \mathbf{E}_{in}^+(\mathbf{r}, t) \\ & - \hbar \sum_{\mathbf{k}\lambda} g_{\mathbf{k}\lambda}^* \hat{c}_{\mathbf{k}\lambda}^\dagger(t) \exp(-i\mathbf{k}\mathbf{r}) \hat{\psi}_1(\mathbf{r}, t) \\ & - \hbar \hat{\psi}_1(\mathbf{r}, t) \sum_{\mathbf{k}\lambda} g_{\mathbf{k}\lambda} \exp(i\mathbf{k}\mathbf{r}) \hat{c}_{\mathbf{k}\lambda}(t), \quad (8) \end{aligned}$$

$$\begin{aligned} i\hbar \frac{\partial \hat{c}_{\mathbf{k}\lambda}(t)}{\partial t} = & \hbar \omega_k \hat{c}_{\mathbf{k}\lambda}(t) - \hbar g_{\mathbf{k}\lambda}^* \int d\mathbf{r} \exp(-i\mathbf{k}\mathbf{r}) \\ & \times \left[\hat{\psi}_2^\dagger(\mathbf{r}, t) \hat{\psi}_1(\mathbf{r}, t) + \hat{\psi}_1^\dagger(\mathbf{r}, t) \hat{\psi}_2(\mathbf{r}, t) \right], \quad (9) \end{aligned}$$

$$g_{\mathbf{k}\lambda} = i \sqrt{\frac{2\pi\omega_k}{\hbar V}} \mathbf{d} \mathbf{e}_\lambda,$$

where $\mathbf{E}_{in}^\pm$ are the positive and negative frequency parts of the incident classical electric field. The operator products in Eqs.(7),(8),(9) are taken in normally ordered form.

The formal solution of (9) for the photon operators is

$$\begin{aligned} \hat{c}_{\mathbf{k}\lambda}(t) = & \hat{c}_{\mathbf{k}\lambda}(0) \exp(-i\omega_k t) \\ & + i g_{\mathbf{k}\lambda}^* \int_0^t dt' \int d\mathbf{r}' \exp[i\omega_k(t' - t) - i\mathbf{k}\mathbf{r}'] \\ & \times \left[\hat{\psi}_2^\dagger(\mathbf{r}', t') \hat{\psi}_1(\mathbf{r}', t') + \hat{\psi}_1^\dagger(\mathbf{r}', t') \hat{\psi}_2(\mathbf{r}', t') \right], \quad (10) \end{aligned}$$

where the first term $\hat{c}_{\mathbf{k}\lambda}(0)$ refers to the free-space photon field and the second one goes back to the interaction with the atoms.

To study the back reaction of the photons on matter we insert (10) in (7) and (8). In the rotating wave approximation we obtain the dynamical equations for the operators of the two matter fields

$$i\hbar \frac{\partial \hat{\psi}_1(\mathbf{r}, t)}{\partial t} = -\frac{\hbar^2 \nabla^2}{2m} \hat{\psi}_1(\mathbf{r}, t) - \mathbf{d} \hat{\mathbf{E}}_{loc}^-(\mathbf{r}, t) \hat{\psi}_2(\mathbf{r}, t), \quad (11)$$

$$\begin{aligned} i\hbar \frac{\partial \hat{\psi}_2(\mathbf{r}, t)}{\partial t} = & -\frac{\hbar^2 \nabla^2}{2m} \hat{\psi}_2(\mathbf{r}, t) + \hbar(\omega_a + \delta - i\gamma/2) \hat{\psi}_2(\mathbf{r}, t) \\ & - \hat{\psi}_1(\mathbf{r}, t) \mathbf{d} \hat{\mathbf{E}}_{loc}^+(\mathbf{r}, t), \quad (12) \end{aligned}$$

where δ and γ are the Lamb shift and the spontaneous emission rate of a single atom in free space, respectively, and we have introduced the operator of the local electric field

$$\begin{aligned} \hat{\mathbf{E}}_{loc}^+(\mathbf{r}, t) = & \mathbf{E}_{in}^+(\mathbf{r}, t) \\ & + i \sum_{\mathbf{k}\lambda} \sqrt{\frac{2\pi\hbar\omega_k}{V}} \mathbf{e}_\lambda \hat{c}_{\mathbf{k}\lambda}(0) \exp(i\mathbf{k}\mathbf{r} - i\omega_k t) \\ & + \int d\mathbf{r}' \nabla \times \nabla \times \frac{\hat{\mathbf{P}}^+(\mathbf{r}', t - R/c)}{R}, \quad (13) \end{aligned}$$

and $\nabla \times$ refers to the point $\mathbf{r}$. Note that in eq.(13) a small volume around the observation point $\mathbf{r}$ is excluded from the integration.

Eq. (13) shows that $\hat{\mathbf{E}}_{loc}^\pm(\mathbf{r}, t)$ is the superposition of the incident field $\mathbf{E}_{in}^\pm(\mathbf{r}, t)$, vacuum fluctuations of the photon field, and the electric field radiated by all other atoms. As one should expect it is this local field, which drives the inner atomic transition in (11), (12).

Equations (11), (12) can be regarded as an atom-optical analogue of the optical Bloch equations [14,11]. They describe the dynamical evolution of the second quantized matter in the field of electromagnetic radiation.

IV. LORENTZ-LORENZ RELATION AND THE SYSTEM OF MAXWELL-BLOCH EQUATIONS IN ATOM OPTICS

A. Local-field correction

Treating the quantum gas as a medium, it is important to establish a connection to the observables of the quantum electrodynamics in media. Having an application to an ultracold gas in mind, we have to relate the local field $\hat{\mathbf{E}}_{loc}(\mathbf{r}, t)$ to the *macroscopic (or mean) field* $\hat{\mathbf{E}}_{mac}(\mathbf{r}, t)$ which is obtained by averaging in space over a region which contains a great number of atoms. For a very low density ultracold gas the region on which the macroscopic field is averaged may contain actually fewer than one atom, yet the averaging technique is still valid due to atomic delocalization.

We generalize the idea of Lorentz as it is described in [11–13] and assume the local field to be a sum of several contributions:

$$\hat{\mathbf{E}}_{loc}(\mathbf{r}, t) = \hat{\mathbf{E}}_{near}(\mathbf{r}, t) + \hat{\mathbf{E}}_{mac}(\mathbf{r}, t) - \hat{\mathbf{E}}_P(\mathbf{r}, t). \quad (14)$$

Let us consider a macroscopically small, but microscopically large volume $\mathcal{V}$ around $\mathbf{r}$. $\hat{\mathbf{E}}_{near}$ is a contribution from the dipoles within $\mathcal{V}$. $\hat{\mathbf{E}}_{mac}$ is the macroscopic field obtained from the field of all dipoles in the medium. Because the dipoles in $\mathcal{V}$ are already taken into account in $\hat{\mathbf{E}}_{near}$, we have to subtract from $\hat{\mathbf{E}}_{mac}$ the contribution $\hat{\mathbf{E}}_P$ of these dipoles as it is obtained in the averaged continuum approximation. $\hat{\mathbf{E}}_P$ is therefore a macroscopic quantity. It can be related to the polarization, which is an averaged quantity too, according to

$$\hat{\mathbf{E}}_P(\mathbf{r}, t) = -\frac{4\pi}{3} \hat{\mathbf{P}}(\mathbf{r}, t). \quad (15)$$

For a novel derivation see [11]. In the following we restrict to the purely classical local-field correction which corresponds to a vanishing $\hat{\mathbf{E}}_{near}$ [11–13]. Thus, we see that, because of the influence of the near dipole-dipole interactions, the local field is obtained from the macroscopic field in adding the *local-field correction* [11–13].

$$\hat{\mathbf{E}}_{loc}^{\pm}(\mathbf{r}, t) = \hat{\mathbf{E}}_{mac}^{\pm}(\mathbf{r}, t) + \frac{4\pi}{3} \hat{\mathbf{P}}^{\pm}(\mathbf{r}, t). \quad (16)$$

This equation is also often called the *Lorentz-Lorenz relation*. The propagation of the operator of the macroscopic field is thereby given by Maxwell wave equation for a charge-free and current-free polarization medium.

B. Nonlinear matter equation

In the next step we decouple the dynamical equations for the two matter field operators. We substitute (16) in (11) and (12) and pass to the reference frame rotating with the frequency ω_L of the incident field which is assumed to be monochromatic

$$\begin{aligned} \hat{\mathbf{E}}_{mac}^+(\mathbf{r}, t) &= \hat{\mathcal{E}}_{mac}^+(\mathbf{r}) \exp(-i\omega_L t), \\ \hat{\psi}_2(\mathbf{r}, t) &= \hat{\phi}_2(\mathbf{r}, t) \exp(-i\omega_L t), \end{aligned} \quad (17)$$

to obtain

$$\begin{aligned} i\hbar \frac{\partial \hat{\psi}_1}{\partial t} &= -\frac{\hbar^2 \nabla^2}{2m} \hat{\psi}_1 - \frac{\hbar}{2} \hat{\Omega}^-(\mathbf{r}) \hat{\phi}_2 - \frac{4\pi}{3} d^2 \hat{\phi}_2^\dagger \hat{\psi}_1 \hat{\phi}_2, \quad (18) \\ i\hbar \frac{\partial \hat{\phi}_2}{\partial t} &= -\frac{\hbar^2 \nabla^2}{2m} \hat{\phi}_2 - \frac{\hbar}{2} \hat{\psi}_1 \hat{\Omega}^+(\mathbf{r}) - \frac{4\pi}{3} d^2 \hat{\psi}_1 \hat{\psi}_1^\dagger \hat{\phi}_2 \\ &\quad - \hbar(\Delta + i\gamma/2) \hat{\phi}_2, \end{aligned} \quad (19)$$

with the detuning $\Delta = \omega_L - \omega_a - \delta$. The position dependent Rabi frequency $\hat{\Omega}^{\pm}(\mathbf{r}) = 2\mathbf{d}\hat{\mathcal{E}}_{mac}^{\pm}(\mathbf{r})/\hbar$ is related to the macroscopic electric field.

We assume sufficiently large detuning such that the contribution of the spontaneous emission is small. We may therefore apply the *adiabatic approximation* to (19), which gives

$$\hat{\phi}_2(\mathbf{r}, t) = -\frac{\hat{\Omega}^+(\mathbf{r}) \hat{\psi}_1(\mathbf{r}, t)}{2[\hat{\Delta}_l(\mathbf{r}, t) + i\gamma/2]}, \quad (20)$$

and eliminate the field $\hat{\psi}_2$ from our scheme. The position dependent *local detuning* introduced in

$$\hat{\Delta}_l(\mathbf{r}, t) = \Delta + \frac{4\pi}{3\hbar} d^2 \hat{\psi}_1^\dagger(\mathbf{r}, t) \hat{\psi}_1(\mathbf{r}, t) \quad (21)$$

is an operator depending on the density operator of the ground state. We interpret this as a shift of the internal energy levels which may increase or decrease with increasing density depending on the sign of the detuning Δ . Analogous frequency shifts can be found in nonlinear optics (see, for instance, [11,15] and references therein).

Because we are mainly interested in atom optical problems and want to study the coherent evolution of the center-of-mass motion of the gas, we shall neglect spontaneous emission. This is valid for situations where the absolute value of the local detuning is much bigger than the spontaneous emission rate $|\Delta_l| \gg \gamma$. In order to do

this we drop in the following the vacuum fluctuations and the spontaneous emission rate γ from our equations.

Then substituting (21) in (18) we obtain as the intended result the nonlinear dynamical equation for the matter field operator $\hat{\psi}_1(\mathbf{r}, t)$

$$i\hbar \frac{\partial \hat{\psi}_1(\mathbf{r}, t)}{\partial t} = \left\{ -\frac{\hbar^2 \nabla^2}{2m} + \frac{\hbar}{4} \frac{\Delta}{\hat{\Delta}_l^2(\mathbf{r}, t)} |\hat{\Omega}^+(\mathbf{r})|^2 \right\} \hat{\psi}_1(\mathbf{r}, t). \quad (22)$$

It couples to the macroscopic electric field via the Rabi frequency $\hat{\Omega}^{\pm}(\mathbf{r})$.

On the assumptions stated in the Section I, equation (22) is general. No approximation apart from the adiabatic approximation and the rotating wave approximation has been made. The dynamical dipole-dipole interaction mediated through the exchange of photons is completely contained. The nonlinearity goes back to the density dependent local detuning $\hat{\Delta}_l(\mathbf{r}, t)$ of eq. (21). For increasing density and positive detuning Δ the local detuning grows and correspondingly the nonlinear term in (22) representing the coupling to the macroscopic electric field becomes smaller. On the other hand, for negative detuning the absolute value of the local detuning decreases with the increase of the density and the nonlinearity becomes greater.

C. Microscopic and macroscopic Maxwell equations

In the multipolar formulation of QED the operator of the microscopic displacement field is given by eq.(5) and the operator of the magnetic field has the form

$$\begin{aligned} \hat{\mathbf{B}}_{mic}(\mathbf{r}, t) &= i \sum_{\mathbf{k}\lambda} \sqrt{\frac{2\pi\hbar\omega_k}{V}} \frac{\omega_k}{c} (\mathbf{k} \times \mathbf{e}_\lambda) \hat{c}_{\mathbf{k}\lambda}(t) \exp(i\mathbf{k}\mathbf{r}) \\ &\quad + H.c. \end{aligned} \quad (23)$$

Calculating time and spatial derivatives of the operators (5), (23) we get microscopic Maxwell equations

$$\begin{aligned} \nabla \times \hat{\mathbf{E}}_{mic} &= -\frac{1}{c} \frac{\partial}{\partial t} \hat{\mathbf{B}}_{mic}, \quad \text{div} \hat{\mathbf{B}}_{mic} = 0, \\ \nabla \times \hat{\mathbf{H}}_{mic} &= \frac{1}{c} \frac{\partial}{\partial t} \hat{\mathbf{D}}_{mic}, \quad \text{div} \hat{\mathbf{D}}_{mic} = 0, \end{aligned} \quad (24)$$

where $\hat{\mathbf{D}}_{mic} = \hat{\mathbf{E}}_{mic} + 4\pi\hat{\mathbf{P}}$ and $\hat{\mathbf{B}}_{mic} = \hat{\mathbf{H}}_{mic}$. The same equations were obtained by G.Juzeliunas [16] in a polariton model of a dielectric medium interacting with light.

In order to get the macroscopic Maxwell equations we have to average eqs.(24) over the microscopically large, but macroscopically small volume of space. However, in the case of ultracold atomic gases due to the atomic delocalization any point of space contains contributions from all the atoms. In this sense the microscopic field is

already averaged and we may simply replace the microscopic fields by their macroscopic analogues.

Assuming that the spatial variations of the atomic density are not rapid and taking into account the adiabatic solution (20), one can rewrite the system of macroscopic Maxwell equations in the form of the wave equation

$$\nabla^2 \hat{\mathcal{E}}_{mac}^\pm + k_L^2 \hat{n}^2 \hat{\mathcal{E}}_{mac}^\pm = 0, \quad (25)$$

where the refractive index is given by the Clausius-Mossotti relation

$$\hat{n}^2 = \frac{1 + \frac{8\pi}{3} \alpha \hat{\psi}_1^\dagger \hat{\psi}_1}{1 - \frac{4\pi}{3} \alpha \hat{\psi}_1^\dagger \hat{\psi}_1}. \quad (26)$$

Eqs.(22),(25) and (26) constitute the general system of Maxwell-Bloch equations, which can be employed for the solution of different problems in nonlinear atom optics. In our paper [9] we have shown that the equations which are usually used for the description of an ultracold ensemble put in an external laser field can be derived from (22),(25) and (26) assuming that the density of atoms is a small parameter.

D. Equal-time commutation relations for the operators of the electromagnetic field

The operators of the local electric and magnetic field strength have the following form:

$$\begin{aligned} \hat{\mathbf{E}}_{loc}(\mathbf{r}, t) &= i \sum_{\mathbf{k}\lambda} \sqrt{\frac{2\pi\hbar\omega_k}{V}} \mathbf{e}_\lambda \hat{c}_{\mathbf{k}\lambda}(t) \exp(i\mathbf{k}\mathbf{r}) + H.c. \\ &\quad - \frac{8\pi}{3} \hat{\mathbf{P}}(\mathbf{r}, t) \end{aligned} \quad (27)$$

$$\begin{aligned} \hat{\mathbf{B}}_{loc}(\mathbf{r}, t) &= i \sum_{\mathbf{k}\lambda} \sqrt{\frac{2\pi\hbar\omega_k}{V}} \frac{\omega_k}{c} (\mathbf{k} \times \mathbf{e}_\lambda) \hat{c}_{\mathbf{k}\lambda}(t) \exp(i\mathbf{k}\mathbf{r}) \\ &\quad + H.c. \end{aligned} \quad (28)$$

Taking into account the commutation relations for the photons and matter-field operators

$$\begin{aligned} [\hat{c}_{\mathbf{k}\lambda}(t), \hat{c}_{\mathbf{k}\lambda}^\dagger(t)] &= 1, \\ [\hat{c}_{\mathbf{k}\lambda}(t), \hat{\psi}_{1,2}(t)] &= [\hat{c}_{\mathbf{k}\lambda}^\dagger(t), \hat{\psi}_{1,2}^\dagger(t)] = 0, \end{aligned} \quad (29)$$

we obtain the following equal-time commutation relations for the operators of the local electromagnetic field

$$\begin{aligned} [\hat{\mathbf{E}}_{loc}^m(\mathbf{r}_1, t), \hat{\mathbf{E}}_{loc}^n(\mathbf{r}_2, t)] &= \left(\frac{8\pi}{3}\right)^2 d^m d^n \delta(\mathbf{r}_1 - \mathbf{r}_2) \\ &\times [\hat{\psi}_1^\dagger(\mathbf{r}_1, t) \hat{\psi}_1(\mathbf{r}_2, t) - \hat{\psi}_1^\dagger(\mathbf{r}_2, t) \hat{\psi}_1(\mathbf{r}_1, t) \\ &\quad + \hat{\psi}_2^\dagger(\mathbf{r}_1, t) \hat{\psi}_2(\mathbf{r}_2, t) - \hat{\psi}_2^\dagger(\mathbf{r}_2, t) \hat{\psi}_2(\mathbf{r}_1, t)], \end{aligned} \quad (30)$$

$$[\hat{\mathbf{E}}_{loc}^m(\mathbf{r}_1, t), \hat{\mathbf{B}}_{loc}^n(\mathbf{r}_2, t)] = -i4\pi\hbar c \sum_l \varepsilon_{mnl} \frac{\partial}{\partial R_l} \delta(\mathbf{R}). \quad (31)$$

In eq.(30) the δ -function vanishes, if $\mathbf{r}_1 \neq \mathbf{r}_2$. If $\mathbf{r}_1 = \mathbf{r}_2$, the term in square brackets in the r.h.s. of eq.(30) vanishes. Therefore, the commutator (30) vanishes for any $\mathbf{r}_1$ and $\mathbf{r}_2$, and we have the canonical commutation relations for the operators of the local electromagnetic field.

Starting from the Lorentz-Lorenz relation (16) and making use of the equal-time commutation relations (30), (31) we get the canonical commutation relations for the operators of the macroscopic electromagnetic field:

$$[\hat{\mathbf{E}}_{mac}^m(\mathbf{r}_1, t), \hat{\mathbf{E}}_{mac}^n(\mathbf{r}_2, t)] = 0, \quad (32)$$

$$\begin{aligned} [\hat{\mathbf{E}}_{mac}^m(\mathbf{r}_1, t), \hat{\mathbf{B}}_{mac}^n(\mathbf{r}_2, t)] &= \\ -i4\pi\hbar c \sum_l \varepsilon_{mnl} \frac{\partial}{\partial R_l} \delta(\mathbf{R}). \end{aligned} \quad (33)$$

The commutation relations (30)-(33) show that our theory is consistent with the general principles of the canonical quantization of the electromagnetic field in a medium.

V. CONCLUSION

We have treated the interaction of dense Bose and Fermi gases with light in the microscopic approach. Our results are not limited to low densities and include the dipole-dipole interaction consistently. The present analysis is concentrated on the common features of the interaction of Bose and Fermi gases with the radiation field. The differences appear when we consider correlation functions which reflect Bose or Fermi statistics. The effects caused by the statistics were considered in [1–3], where it was shown that the refractive index contains additional terms. However, these statistical effects are not very important because the leading term is governed by the Clausius-Mossotti formula.

ACKNOWLEDGMENTS

This work has been supported by the Deutsche Forschungsgemeinschaft and the Optikzentrum Konstanz. One of us (K.V.K.) would like to thank the members of the AG Audretsch at the University of Konstanz for many interesting discussions and kind hospitality. Valuable discussions with G.Hegerfeldt, D.-G.Welsch, S.Scheel, G.Juzeliunas, J.Ruostekoski, and the group of Prof.W.Schleich at Ulm university are gratefully acknowledged. Special thanks to C.M.Bowden for his interest to our work.

- [1] Morice, O., Castin, Y., and Dalibard, J., 1995, *Phys.Rev.A*, **51**, 3896.
- [2] Ruostekoski, J., and Javanainen, J., 1997, *Phys.Rev.A*, **55**, 513.
- [3] Ruostekoski, J., and Javanainen, J., 1999, *cond-mat/9902172*.
- [4] Lenz, G., Meystre, P., and Wright, E., 1993, *Phys.Rev.Lett.*, **71**, 3271.
- [5] Zhang, W., and Walls, D., 1994, *Phys.Rev.A*, **49**, 3799.
- [6] Lenz, G., Meystre, P., and Wright, E., 1994, *Phys.Rev.A*, **50**, 1681.
- [7] Castin, Y., and Mølmer, K., 1995, *Phys.Rev.A*, **51**, R3426.
- [8] Wallis, H., 1997, *Phys.Rev.A*, **56**, 2060.
- [9] Krutitsky, K.V., Burgbacher, F., and Audretsch, J., 1999, *Phys.Rev.A*, **59**, 1517.
- [10] Cohen-Tannoudji, C., Dupont-Roc, J., and Grynberg, G., 1989, *Photons and Atoms: Introduction to Quantum Electrodynamics* (New York: Wiley & Sons).
- [11] Bowden, C.M., and Dowling, J.P., 1993, *Phys.Rev.A*, **47**, 1247.
- [12] Born, M., and Wolf, E., 1970, *Principles of Optics* (New York: Pergamon).
- [13] Jackson, J., 1975, *Classical Electrodynamics* (New York: Wiley).
- [14] Allen, L., and Eberly, J., 1978, *Optical Resonance and Two-Level Atoms* (New York: Wiley).
- [15] Crenshaw, M.E., and Bowden, C.M., 1996, *Phys.Rev.A*, **53**, 1139.
- [16] Juzeliunas, G., 1997, *Phys.Rev.A*, **55**, 929.